

AN EXACT TEST FOR THE CHOICE OF THE COMBINATION OF FIRST  
DIFFERENCES AND PERCENTAGE CHANGES IN LINEAR MODELS

Wai Cheung Ip

Department of Applied Mathematics

The Hong Kong Polytechnic University

Hung Hom

Kowloon

Hong Kong

Email: mathipwc@polyu.edu.hk

**Abstract**

Econometric models are often formulated in terms of the first difference or the percentage change, generalized by the Box-Cox difference transformation. The choice of suitable functional forms has relied heavily upon established statistical procedures which adopt primarily the likelihood approach and confine to a single transformation parameter only. We have derived an exact test for the parameter vector of transformation in linear models. By utilizing Taylor series approximations this reduces to a choice between two regression equations. The test statistic which has an exact  $F$ -distribution can be easily calculated from these two regressions by least squares algorithm. Monte Carlo results demonstrate that our proposed test is more capable than the likelihood approach in capturing the correct size yet is as powerful as the latter. It is therefore a simple and ready statistical procedure for assessing the suitable choice of the combination of first differences and percentage changes in economic forecast models.

JEL Classification: C12

Key Words: functional form, Box-Cox Transformation, Taylor series, least squares

## 1. Introduction

The Box-Cox transformation of a positive series  $\{ y_t \}$  is given by

$$(1.1) \quad y_t^{(I)} = \begin{cases} \frac{y_t^I - 1}{I}, & I \neq 0, \\ \ln y_t, & I = 0 \end{cases}, \quad t = 1, \dots, T$$

where  $y_t$  is an economic variable at time period  $t$  and  $I$  is a Box-Cox transformation parameter. Typically many econometric applications examine not the level of the variable, but its change per time period. Examples include the St. Louis equation due to Anderson and Carlson (1970), the money demand model by Hafer and Hein (1980) and the consumer index of Colclough and Lange (1982). A discrete approximation to the time derivative of  $y_t^{(I)}$  is given by Layson and Seaks (1984) as

$$(1.2) \quad \Delta y_t^{(I)} = y_{t-1}^{I-1} \Delta y_t, \quad t = 1, \dots, T,$$

where  $\Delta y_t = y_t - y_{t-1}$  denotes the first difference. The usefulness of (1.2) is that  $I=1$  yields the first difference while  $I=0$  gives the percentage change. This is termed the Box-Cox difference transformation by Seaks and Vines (1990). Both the first difference and the percentage change of an economic variable may therefore be generalized by  $\Delta y_t^{(I)}$ . Thus a regression model which utilizes both the first difference and the percentage change can be represented by

$$(1.3) \quad \Delta y_t^{(I_1)} = \mathbf{a}_1 x_{1t} + \dots + \mathbf{a}_p x_{pt} + \Delta z_{2t}^{(I_2)} \mathbf{b}_2 + \dots + \Delta z_{qt}^{(I_q)} \mathbf{b}_q + \mathbf{e}_t,$$

where  $I_i$  ( $i=1, \dots, q$ ) are Box-Cox difference transformation parameters,  $x_{jt}$  ( $j=1, 2, \dots, p$ ) are observations on the  $p$  independent variables which are not transformed,  $\Delta y_t^{(I_1)}$  is defined as in (1.2),  $\Delta z_{it}^{(I_i)}$ , ( $i=2, \dots, q$ ) are similarly defined,  $\mathbf{a}_j$  ( $j=1, \dots, p$ ) and  $\mathbf{b}_i$  ( $i=2, \dots, q$ ) are regression coefficients, and  $\mathbf{e}_t$  is the error term. We may regard (1.3) as a general model in which the

transformed parameters permit more general forms than first differences or percentage changes though  $I = 0$  or 1 has a straightforward and simple interpretation.

Unfortunately model selection is a very difficult task in practice. As Dhrymes et al. (1972, p294) have pointed out that economic theory typically provides little guidance as to the proper functional form appropriate to the specification of the economic relationships. The choice of suitable functional forms in econometric applications thus relies heavily upon established statistical procedures.

In this paper, we shall develop an Andrews (1971) type procedure for testing the choice of the set of Box-Cox difference transformation parameters in (1.3), allowing different transformation parameters for different variables. Its statistical methodology follows the line of Milliken and Graybill (1970) and is based on the  $F$ -distribution. Section 2 reviews existing single transformation parameter procedures. In section 3 we shall develop the testing procedure which reduces to a choice between two regression equations. The alternative regression, though of no economic interest, is easy to understand. Section 4 outlines the steps how the test statistic can be computed using a standard statistical package. In section 5 the new procedure is compared with the likelihood ratio approach by Monte Carlo models. Section 6 presents the conclusions of this paper.

## **2. Single Transformation Parameter Procedures**

Existing procedures include the likelihood ratio (LR) test and variants of the Lagrange multiplier (LM) tests. They assume the same transformation parameter be used on both the dependent and independent variables. The procedure of Layson and Seaks (1984) is primarily a LR test. It requires a search for an optimal solution of the unrestricted model which in practice can be completed by a sequence of regressions on transformed data with various values of the transformation parameter over a grid. Though the first difference of the dependent variable and percentage change of the regressor are used in the poverty and growth model of Thornton et al. (1978) Layson and Seaks (1984) have only applied their procedure to test the latter choice but not the use of the first difference of the dependent variable. The search could be very expensive if not impossible when different parameters of transformation are allowed. Those of Coulson

and Robins (1985) and of Park (1991) are different variants of LM tests, constructed from rather complicated auxiliary regressions in which the regressors involve either the time derivatives of the transformed variables or the derivatives of the unrestricted log-likelihood function with respect to the individual variables. The regressions are artificially constructed to generate tests for misspecification and their relations to the hypothesis to be tested are not obvious. The computation involved is by no means simple or straightforward. Further, the LR and LM procedures are asymptotic and therefore are only approximate in small and moderate sample situations. The test statistic of the proposed procedure can be calculated easily by ordinary least squares estimation of two regression equations, thus avoiding the relatively more laborious search over a grid of values.

### 3. The Test Procedure

Using matrix notation we may express (1.3) as

$$(3.1) \quad \Delta \mathbf{y}^{(I_1)} = \mathbf{X}\mathbf{a} + \Delta \mathbf{Z}^{(\mathbf{I}_2)}\mathbf{b} + \mathbf{e}$$

where  $\mathbf{I}_2=(I_2 \ I_3 \ \dots \ I_q)'$ ,  $\Delta \mathbf{y}^{(I_1)}$  is the  $T \times 1$  vector of transformed dependent variables,  $\mathbf{X}$  is the  $T \times p$  matrix of original independent variables,  $\Delta \mathbf{Z}^{(\mathbf{I}_2)} = (\Delta \mathbf{z}_2^{(I_2)} \ \Delta \mathbf{z}_3^{(I_3)} \ \dots \ \Delta \mathbf{z}_q^{(I_q)})$  is the  $T \times (q-1)$  matrix of transformed independent variables,  $\mathbf{a}=(\mathbf{a}_1 \ \mathbf{a}_2 \ \dots \ \mathbf{a}_p)'$  is a  $p \times 1$  and  $\mathbf{b}=(\mathbf{b}_2 \ \mathbf{b}_3 \ \dots \ \mathbf{b}_q)'$  is a  $(q-1) \times 1$  vectors of coefficients, and  $\mathbf{e}$  is the  $T \times 1$  vector of independently and normally distributed disturbances with zero mean and constant variance  $\sigma^2$ . Thus the model selection problem reduces to testing the hypothesis about the transformation parameter vector  $\mathbf{I}=(I_1 \ I_2 \ \dots \ I_q)'$  in model (3.1).

Expanding  $\Delta \mathbf{y}^{(I_1)}$  in Taylor series about a hypothetical value  $\mathbf{I}_1^{(0)}$  yields

$$(3.2) \quad \Delta \mathbf{y}^{(I_1)} \approx \Delta \mathbf{y}^{(I_1^{(0)})} + (\mathbf{I}_1 - \mathbf{I}_1^{(0)}) \mathbf{w}(\mathbf{I}_1^{(0)})$$

where  $\mathbf{w}(\mathbf{I}_1^{(0)}) = \partial \Delta \mathbf{y}^{(I_1)} / \partial \mathbf{I}_1$  evaluated at  $\mathbf{I}_1 = \mathbf{I}_1^{(0)}$ . It can be readily shown that  $\mathbf{w}(\mathbf{I}_1^{(0)})$  has  $t$ -th element

$$(3.3) \quad w_t(\mathbf{I}_1^{(0)}) = (\ln y_{t-1}) \Delta y_t^{(I_1^{(0)})}.$$

Similarly, we have

$$\Delta \mathbf{Z}^{(\mathbf{I}_2)} \approx \Delta \mathbf{Z}^{(\mathbf{I}_2^{(0)})} + \mathbf{V}(\mathbf{I}_2^{(0)}) (\mathbf{I}_2 - \mathbf{I}_2^{(0)})$$

where  $\mathbf{I}_2^{(0)} = (\mathbf{I}_2^{(0)} \ \mathbf{I}_3^{(0)} \ \dots \ \mathbf{I}_q^{(0)})'$  is a hypothetical value of  $\mathbf{I}_2$ ,

$$(3.4) \quad \mathbf{V}(\mathbf{I}_2^{(0)}) = [\mathbf{v}_2(\mathbf{I}_2^{(0)}) \ \mathbf{v}_3(\mathbf{I}_3^{(0)}) \ \dots \ \mathbf{v}_q(\mathbf{I}_q^{(0)})]$$

and  $\mathbf{v}_i(\mathbf{I}_i^{(0)}) = \partial \Delta \mathbf{Z}^{(I_i)} / \partial \mathbf{I}_i$  being evaluated at  $\mathbf{I}_i = \mathbf{I}_i^{(0)}$ ,  $i=2, \dots, q$ . The individual elements of  $\mathbf{v}_i(\mathbf{I}_i^{(0)})$  are given similarly to (3.3). Substituting (3.2) and (3.4) into (3.1) and writing  $\mathbf{I}_0 = (\mathbf{I}_1^{(0)} \ \mathbf{I}_2^{(0)} \ \dots \ \mathbf{I}_q^{(0)})'$ , we shall obtain

$$(3.5) \quad \Delta \mathbf{y}^{(I_1^{(0)})} = \mathbf{X} \mathbf{a} + \Delta \mathbf{Z}^{(\mathbf{I}_2^{(0)})} \mathbf{b} + [-\mathbf{w}(\mathbf{I}_1^{(0)}) : \mathbf{V}(\mathbf{I}_2^{(0)}) \mathbf{b}] (\mathbf{I} - \mathbf{I}_0) + \mathbf{e}$$

which, on letting  $\mathbf{g} = \mathbf{I} - \mathbf{I}_0$  and  $\mathbf{U}(\mathbf{I}_0, \mathbf{b}) = [-\mathbf{w}(\mathbf{I}_1^{(0)}) : \mathbf{V}(\mathbf{I}_2^{(0)}) \mathbf{b}]$ , may be rewritten as

$$(3.6) \quad \Delta \mathbf{y}^{(I_1^{(0)})} = \mathbf{X}\mathbf{a} + \Delta \mathbf{Z}^{(I_2^{(0)})}\mathbf{b} + \mathbf{U}(\mathbf{I}_0, \mathbf{b})\mathbf{g} + \mathbf{e}.$$

Box (1980) calls the new explanatory variable  $\mathbf{U}$  a *constructed variable*. The testing of the null hypothesis  $H_0: \mathbf{I} = \mathbf{I}_0$  against the alternative  $H_1: \mathbf{I} \neq \mathbf{I}_0$  has become a choice between (3.6) and the null model

$$(3.7) \quad \Delta \mathbf{y}^{(I_1^{(0)})} = \mathbf{X}\mathbf{a} + \Delta \mathbf{Z}^{(I_2^{(0)})}\mathbf{b} + \mathbf{e}.$$

Equivalently, we are testing  $H_0: \mathbf{g} = \mathbf{0}$  against  $H_1: \mathbf{g} \neq \mathbf{0}$  in model (3.6). To eliminate the dependence of  $\mathbf{U}$  on  $\mathbf{e}$  in (3.6) the former will be replaced by its least squares estimate from the null model. Let  $\hat{\mathbf{a}}$  and  $\hat{\mathbf{b}}$  be the least squares estimates of  $\mathbf{a}$  and  $\mathbf{b}$  in (3.7), the fitted values  $\Delta \hat{y}_t^{(I_1^{(0)})}$  of  $\Delta y_t^{(I_1^{(0)})}$  can be computed from (3.7) after replacing the regression coefficients by their least squares estimates, and the fitted values of  $y_t^{(I_1^{(0)})}$  can thus be determined recursively by

$$(3.8) \quad \begin{aligned} \hat{y}_1^{(I_1^{(0)})} &= \Delta \hat{y}_1^{(I_1^{(0)})} + y_0^{(I_1^{(0)})} \\ \hat{y}_2^{(I_1^{(0)})} &= \Delta \hat{y}_2^{(I_1^{(0)})} + \hat{y}_1^{(I_1^{(0)})} \\ &\vdots \\ \hat{y}_T^{(I_1^{(0)})} &= \Delta \hat{y}_T^{(I_1^{(0)})} + \hat{y}_{T-1}^{(I_1^{(0)})} \end{aligned}$$

By the inverse of the Box-Cox formula (1.1) the fitted values of  $y_t$  are given by

$$(3.9) \quad \hat{y}_t = \begin{cases} (\mathbf{I}_1^{(0)} \hat{y}_t^{(I_1^{(0)})} + 1)^{1/I_1^{(0)}}, & \mathbf{I}_1^{(0)} \neq 0 \\ e^{\hat{y}_t^{(0)}}, & \mathbf{I}_1^{(0)} = 0 \end{cases}, t = 1, \dots, T$$

Substituting  $\Delta \hat{y}_t^{(I_1^{(0)})}$  and  $\ln \hat{y}_{t-1}$  into (3.3) we shall get  $\hat{w}_t(I_1^{(0)})$  and hence  $\hat{\mathbf{w}}(I_1^{(0)})$ , the least squares estimate of  $\mathbf{w}(I_1^{(0)})$ , and finally

$$(3.10) \quad \hat{\mathbf{U}} = [-\hat{\mathbf{w}}(I_1^{(0)}) : \mathbf{V}(\mathbf{I}_2^{(0)}) \hat{\mathbf{b}}]$$

upon substituting  $\hat{\mathbf{b}}$  in  $\mathbf{U}(\mathbf{I}_0, \mathbf{b})$ . Replacing  $\mathbf{U}$  in (3.6) by  $\hat{\mathbf{U}}$  gives

$$(3.11) \quad \Delta \mathbf{y}^{(I_1^{(0)})} = \mathbf{X}\mathbf{a} + \Delta \mathbf{Z}^{(I_2^{(0)})} \mathbf{b} + \hat{\mathbf{U}}\mathbf{g} + \mathbf{e}$$

which satisfies the standard conditions of ordinary least squares. The use of  $\hat{\mathbf{U}}$  to replace the unobservable  $\mathbf{U}$  has been put forward by Milliken and Graybill (1970) for two obvious reasons, namely,  $\hat{\mathbf{U}}$  can be computed from the data and is independent of the disturbance. The same tactic has also been used in Andrews (1971). Let  $S_0$  and  $S_1$  be the residual sums of squares by least squares on the regression equations (3.7) and (3.11) respectively. It follows from the results of Milliken and Graybill (1970) that the quantity

$$(3.12) \quad F = \frac{(S_0 - S_1)/q}{S_1/(T - p - 2q + 1)}$$

will follow a  $F$ -distribution with  $q$  and  $T-p-2q+1$  degrees of freedom. Hence  $F$  is a test of the hypothesis  $\mathbf{g}=\mathbf{0}$  or  $\mathbf{l}=\mathbf{l}_0$ .

A detailed development of the above results can be found in Milliken and Graybill (1970) who have shown that  $F$  has a  $F$ -distribution when  $H_0: \mathbf{g}=\mathbf{0}$  is true. The test is 'exact' in the sense that an 'exact' significance will be obtained from which 'exact' confidence limits may be calculated when  $H_0$  is true. However, little is known about  $F$  when  $\mathbf{g} \neq \mathbf{0}$ . See Ward and Dick (1952). Andrews (1971) has pointed out that the precision in (3.11) may affect the efficiency of the test but it will not affect the 'exactness' of the distribution of the test statistic.

The proposed procedure is therefore more capable of capturing the correct size of the test than any other asymptotic ones. For more discussion of other advantages of the proposed procedure over asymptotic ones; see Andrews (1971).

#### 4. Computing Procedures

This section briefly outlines procedures for computing the test statistic  $F$ . The procedures described here assume the use of a statistical package.

The steps are as follows:

1. Transform  $\mathbf{y}$  and  $\mathbf{Z}$  as in (1.2) to get  $\Delta\mathbf{y}^{(I_1^{(0)})}$  and  $\Delta\mathbf{Z}^{(I_2^{(0)})}$  and to form the regression equation (3.7).
2. Estimate equation (3.7) by least squares to obtain coefficient estimates  $\hat{\mathbf{a}}$  and  $\hat{\mathbf{b}}$ , and the error variance estimate  $s_0^2$ , say.
3. Compute  $\Delta\mathbf{y}^{(I_1^{(0)})} = \mathbf{X}\hat{\mathbf{a}} + \Delta\mathbf{Z}^{(I_2^{(0)})}\hat{\mathbf{b}}$ , from which calculate  $\hat{y}_t^{(I_1^{(0)})}$  using the recursive relations (3.8) and  $\hat{y}_t$  by the inverse Box-Cox formula (3.9).
4. Compute  $\hat{w}_t(I_1^{(0)})$  by (3.3) using computed values from step 3 and form the vector  $\hat{\mathbf{w}}(I_1^{(0)})$ . Compute  $\mathbf{V}(I_2^{(0)})$  using a formula similar to (3.3). Augment  $-\hat{\mathbf{w}}(I_1^{(0)})$  and  $\mathbf{V}(I_2^{(0)})\hat{\mathbf{b}}$  to form  $\hat{\mathbf{U}}$  given in (3.10).
5. Form the regression equation (3.11) and estimate it by least squares to obtain error variance estimate  $s_1^2$ , say.
6. Finally calculate the test statistic  $F$  in (3.12) by putting  $S_0=(T-p-q+1)s_0^2$  and  $S_1=(T-p-2q+1)s_1^2$ .

When the package such as RATS used has built-in testing procedure for the regression coefficient vector, Steps 5 and 6 would be combined to one of testing  $\mathbf{g}=\mathbf{0}$  in the regression equation (3.11). The null model is only used to compute the fitted values  $\hat{\mathbf{U}}$  of the alternative model.



## 5. The Monte Carlo

To evaluate our proposed F-test and to compare its performance with that of the LR procedure programmes in RATS are written to generate data for models with combinations of first differences and percentage changes. Sample sizes of 20, 30, 40, 60 and 100 are selected so that both the small sample and asymptotic properties can be studied. Each experiment involves 2000 replications. The four models studied are:

$$\begin{aligned} \text{M00: } \% \Delta y_t &= 0.01 + 0.9 \% \Delta z_t + \mathbf{e}_t \\ \% \Delta z_t &\sim \text{U}(0, 0.07), \quad \mathbf{e}_t \sim \text{N}(0, 0.015^2), \quad z_0 = 1.0, y_0 = 5.0 \end{aligned}$$

$$\begin{aligned} \text{M01: } \% \Delta y_t &= 0.01 + 0.03 \Delta z_t + \mathbf{e}_t \\ \Delta z_t &\sim \text{U}(-0.5, 1.5), \quad \mathbf{e}_t \sim \text{N}(0, 0.015^2), \quad z_0 = 5.0, y_0 = 15.0 \end{aligned}$$

$$\begin{aligned} \text{M10: } \Delta y_t &= 100 + 55 \% \Delta z_t + \mathbf{e}_t \\ \% \Delta z_t &\sim \text{U}(0, 0.07), \quad \mathbf{e}_t \sim \text{N}(0, 1), \quad z_0 = 10, y_0 = 100 \end{aligned}$$

$$\begin{aligned} \text{M11: } \Delta y_t &= 10 + 2 \Delta z_t + \mathbf{e}_t \\ \Delta z_t &\sim \text{U}(0, 2), \quad \mathbf{e}_t \sim \text{N}(0, 1), \quad z_0 = 10, y_0 = 100 \end{aligned}$$

The population  $R^2$  are .593, .571, .553 and .571 respectively. M00 and M11 have been used by Seaks and Vines (1990). The models are tested under each of the null hypotheses:  $H_0: (I_1=0, I_2=0)$ ,  $H_0: (I_1=0, I_2=1)$ ,  $H_0: (I_1=1, I_2=0)$  and  $H_0: (I_1=1, I_2=1)$  in turn, at 1%, 5% and 10% significance levels. The empirical significance level is recorded, when the null hypothesis is true this is the Type I error and when the null is false this will be the power of the relevant test.

A slight modification of Layson and Seak's (1984) result gives the log likelihood function under normality:

$$l = -\frac{T}{2} [\ln(2\mathbf{p}) + \ln \mathbf{s}^2] - \frac{1}{2\mathbf{s}^2} \sum_{t=1}^T \mathbf{e}_t + (\mathbf{I}_1 - 1) \sum_{t=1}^T \ln y_{t-1}.$$

The RATS procedure MAXIMIZE is used to compute the maximum likelihood estimate  $\hat{\mathbf{I}}$  of  $\mathbf{I}$  and to calculate the LR test statistic

$$c^2 = 2[l(\hat{\mathbf{I}}) - l(\mathbf{I}_0)] \sim c_q^2.$$

While our F procedure took few seconds the LR procedure took several minutes to few hours to complete 2000 replications even though the true parameter values were used to start with. When values other than the parameter values were used as initial values the latter procedure often located the optimal solution after several hundred iterations.

Table 1a

Size/Power of Test at 1% Significance Level Nominal

True Model:  $\mathbf{I}_1=0, \mathbf{I}_2=0$

| Null Hypothesis | $\mathbf{I}_1=0, \mathbf{I}_2=0$ |     | $\mathbf{I}_1=0, \mathbf{I}_2=1$ |       | $\mathbf{I}_1=1, \mathbf{I}_2=0$ |       | $\mathbf{I}_1=1, \mathbf{I}_2=1$ |      |       |
|-----------------|----------------------------------|-----|----------------------------------|-------|----------------------------------|-------|----------------------------------|------|-------|
|                 | F                                | LR  | F                                | LR    | F                                | LR    | F                                | LR   |       |
| Sample          | 20                               | 0.8 | 2.1                              | 7.0   | 14.3                             | 34.1  | 31.4                             | 3.0  | 5.4   |
|                 | 30                               | 1.0 | 1.4                              | 42.6  | 58.7                             | 95.2  | 95.5                             | 10.8 | 29.8  |
|                 | 40                               | 0.9 | 1.1                              | 89.6  | 96.2                             | 100.0 | 100.0                            | 23.4 | 79.2  |
| Size            | 60                               | 0.8 | 0.9                              | 100.0 | 100.0                            | 100.0 | 100.0                            | 41.0 | 100.0 |
|                 | 100                              | 0.9 | 0.8                              | 100.0 | 100.0                            | 100.0 | 100.0                            | 65.1 | 100.0 |

Table 1b

Size/Power of Test at 5% Significance Level Nominal

True Model:  $I_1=0, \lambda_2=0$

| Null Hypothesis |     | $I_1=0, I_2=0$ |     | $I_1=0, I_2=1$ |       | $I_1=1, I_2=0$ |       | $I_1=1, I_2=1$ |       |
|-----------------|-----|----------------|-----|----------------|-------|----------------|-------|----------------|-------|
| Test            |     | F              | LR  | F              | LR    | F              | LR    | F              | LR    |
| Sample          | 20  | 5.0            | 4.8 | 23.3           | 28.1  | 60.7           | 52.0  | 12.4           | 12.9  |
|                 | 30  | 5.6            | 4.3 | 70.6           | 76.0  | 98.7           | 98.7  | 27.0           | 50.3  |
|                 | 40  | 4.9            | 3.8 | 97.5           | 99.0  | 100.0          | 100.0 | 44.1           | 91.2  |
| Size            | 60  | 5.0            | 3.7 | 100.0          | 100.0 | 100.0          | 100.0 | 64.6           | 100.0 |
|                 | 100 | 5.0            | 4.5 | 100.0          | 100.0 | 100.0          | 100.0 | 79.6           | 100.0 |

Table 1c

Size/Power of Test at 10% Significance Level Nominal

True Model:  $I_1=0, I_2=0$

| Null Hypothesis |     | $I_1=0, I_2=0$ |     | $I_1=0, I_2=1$ |       | $I_1=1, I_2=0$ |       | $I_1=1, I_2=1$ |       |
|-----------------|-----|----------------|-----|----------------|-------|----------------|-------|----------------|-------|
| Test            |     | F              | LR  | F              | LR    | F              | LR    | F              | LR    |
| Sample          | 20  | 9.4            | 7.9 | 35.3           | 36.6  | 72.5           | 63.8  | 21.1           | 20.0  |
|                 | 30  | 10.5           | 7.3 | 81.5           | 84.1  | 99.5           | 99.4  | 38.0           | 60.7  |
|                 | 40  | 9.9            | 7.2 | 98.8           | 99.5  | 100.0          | 100.0 | 57.0           | 94.7  |
| Size            | 60  | 10.0           | 7.9 | 100.0          | 100.0 | 100.0          | 100.0 | 74.2           | 100.0 |
|                 | 100 | 9.6            | 8.9 | 100.0          | 100.0 | 100.0          | 100.0 | 85.3           | 100.0 |

Tables 1a-1c give the size and power of the tests for model M00. The F-test appears generally to capture the correct size while the LR test overstates the Type I error at the 1% level in small samples and understates it at the 10% level for all sample sizes. It does not appear to give the correct size even for sample sizes as large as 100. The two procedures perform equally well under  $H_0: (I_1=0, I_2=1)$  and  $H_0: (I_1=1, I_2=0)$ . They both lead to correct decision by rejecting the incorrect nulls  $H_0: (I_1=0, I_2=1)$  when  $T \geq 60$  and  $H_0: (I_1=1, I_2=0)$  when  $T \geq 40$  in all cases. When testing under  $H_0: (I_1=1, I_2=1)$  F appears less powerful than LR in rejecting the incorrect null.

Table 2a

Size/Power of Test at 1% Significance Level Nominal

True Model:  $I_1=0, I_2=1$

| Null Hypothesis | $I_1=0, I_2=0$ |       | $I_1=0, I_2=1$ |     | $I_1=1, I_2=0$ |       | $I_1=1, I_2=1$ |       |       |
|-----------------|----------------|-------|----------------|-----|----------------|-------|----------------|-------|-------|
| Test            | F              | LR    | F              | LR  | F              | LR    | F              | LR    |       |
| Sample Size     | 20             | 10.8  | 12.1           | 0.8 | 1.6            | 29.2  | 34.8           | 3.1   | 3.6   |
|                 | 30             | 52.2  | 55.9           | 1.0 | 1.3            | 94.0  | 96.0           | 26.0  | 38.6  |
|                 | 40             | 85.8  | 99.2           | 1.0 | 1.6            | 100.0 | 100.0          | 67.3  | 100.0 |
| Sample Size     | 60             | 98.9  | 100.0          | 0.9 | 0.8            | 100.0 | 100.0          | 98.2  | 100.0 |
|                 | 100            | 100.0 | 100.0          | 0.9 | 1.0            | 100.0 | 100.0          | 100.0 | 100.0 |

Table 2b

Size/Power of Test at 5% Significance Level Nominal

True Model:  $I_1=0, I_2=1$

| Null Hypothesis |     | $I_1=0, I_2=0$ |       | $I_1=0, I_2=1$ |     | $I_1=1, I_2=0$ |       | $I_1=1, I_2=1$ |       |
|-----------------|-----|----------------|-------|----------------|-----|----------------|-------|----------------|-------|
| Test            |     | F              | LR    | F              | LR  | F              | LR    | F              | LR    |
| Sample          | 20  | 28.1           | 25.1  | 4.9            | 2.8 | 57.1           | 54.4  | 11.9           | 11.0  |
|                 | 30  | 77.1           | 76.7  | 4.7            | 3.8 | 99.4           | 98.9  | 48.4           | 47.6  |
|                 | 40  | 96.0           | 100.0 | 5.0            | 4.7 | 100.0          | 100.0 | 84.7           | 100.0 |
| Size            | 60  | 99.8           | 100.0 | 4.2            | 4.9 | 100.0          | 100.0 | 100.0          | 100.0 |
|                 | 100 | 100.0          | 100.0 | 4.7            | 4.1 | 100.0          | 100.0 | 100.0          | 100.0 |

Table 2c

Size/Power of Test at 10% Significance Level Nominal

True Model:  $I_1=0, I_2=1$

| Null Hypothesis |     | $I_1=0, I_2=0$ |       | $I_1=0, I_2=1$ |     | $I_1=1, I_2=0$ |       | $I_1=1, I_2=1$ |       |
|-----------------|-----|----------------|-------|----------------|-----|----------------|-------|----------------|-------|
| Test            |     | F              | LR    | F              | LR  | F              | LR    | F              | LR    |
| Sample          | 20  | 42.1           | 34.9  | 9.5            | 7.9 | 71.1           | 66.7  | 20.7           | 18.1  |
|                 | 30  | 86.9           | 98.4  | 9.9            | 7.3 | 99.8           | 99.4  | 61.3           | 58.6  |
|                 | 40  | 98.6           | 100.0 | 9.7            | 8.0 | 100.0          | 100.0 | 91.3           | 100.0 |
| Size            | 60  | 99.9           | 100.0 | 9.0            | 6.9 | 100.0          | 100.0 | 100.0          | 100.0 |
|                 | 100 | 100.0          | 100.0 | 9.9            | 7.8 | 100.0          | 100.0 | 100.0          | 100.0 |

The size and power for model M01 are given in Tables 2a-2c. The F procedure is seen again to state the correct nominal size. On the other hand, LR tends to overstate the size at the 1% level in small samples and overstate it at the 5 and 10% levels for all sample sizes considered. The Type I error of LR by no means appear to converge to the nominal one even when the sample size is 100. When testing under incorrect hypotheses, both test procedures seem to perform equally well in rejecting the incorrect null. In some cases, a sample size of 40 is sufficiently large to lead to rejection of wrong nulls in 100% of the times.

Table 3a

Size/Power of Test at 1% Significance Level Nominal

True Model:  $I_1=1, I_2=0$

| Null Hypothesis |     | $I_1 = 1, I_2 = 0$ |     | $I_1 = 1, I_2 = 1$ |       |
|-----------------|-----|--------------------|-----|--------------------|-------|
| Test            |     | F                  | LR  | F                  | LR    |
| Sample Size     | 20  | 0.7                | 1.9 | 3.7                | 4.4   |
|                 | 30  | 0.8                | 1.3 | 21.7               | 33.5  |
|                 | 40  | 1.0                | 1.6 | 61.8               | 78.5  |
|                 | 60  | 0.9                | 1.1 | 99.5               | 100.0 |
|                 | 100 | 0.9                | 1.0 | 100.0              | 100.0 |

Table 3b

Size/Power of Test at 5% Significance Level Nominal

True Model:  $I_1=1, I_2=0$

| Null Hypothesis |     | $I_1 = 1, I_2 = 0$ |     | $I_1 = 1, I_2 = 1$ |       |
|-----------------|-----|--------------------|-----|--------------------|-------|
| Test            |     | F                  | LR  | F                  | LR    |
| Sample          | 20  | 4.6                | 4.8 | 15.3               | 10.5  |
|                 | 30  | 4.7                | 3.8 | 46.1               | 52.0  |
|                 | 40  | 4.7                | 4.8 | 83.7               | 90.4  |
| Size            | 60  | 5.2                | 4.6 | 100.0              | 100.0 |
|                 | 100 | 5.1                | 4.5 | 100.0              | 100.0 |

Table 3c

Size/Power of Test at 10% Significance Level Nominal

True Model:  $I_1=1, I_2=0$

| Null Hypothesis |     | $I_1 = 1, I_2 = 0$ |     | $I_1 = 1, I_2 = 1$ |       |
|-----------------|-----|--------------------|-----|--------------------|-------|
| Test            |     | F                  | LR  | F                  | LR    |
| Sample          | 20  | 10.2               | 7.8 | 25.5               | 16.5  |
|                 | 30  | 9.6                | 7.3 | 58.8               | 61.3  |
|                 | 40  | 9.5                | 8.7 | 91.3               | 94.0  |
| Size            | 60  | 10.3               | 7.9 | 100.0              | 100.0 |
|                 | 100 | 9.9                | 9.0 | 100.0              | 100.0 |

For model M10 the two tests are equally powerful when in fact the null were false. When the null is  $H_0: (I_1=0, I_2=0)$  or  $H_0: (I_1=0, I_2=1)$  both of them reject the incorrect null in all cases and therefore results are not presented here. As visible from Tables 3a-3c they do not appear to outperform the other under the incorrect null  $H_0: (I_1=1, I_2=1)$ . They only differ in stating the size under the true null. F generally gives correct sizes at all levels while LR overstates or understates it in most cases.

From Tables 4a-4c it can be seen that F yields Type I errors that are very close to the nominal ones. In contrast LR only produces close to stated sizes in large samples. For example, its type I error is at least two times the nominal value when  $T \leq 40$  at the 1 % level. When the null is  $H_0: (I_1=1, I_2=0)$  both tests do not appear differently in their ability of rejecting the null. When the null is  $H_0: (I_1=0, I_2=0)$  or  $H_0: (I_1=0, I_2=1)$ , both approaches reject the wrong null in all of the cases.

Table 4a  
Size/Power of Test at 1% Significance Level Nominal  
True Model:  $I_1=1, I_2=1$

| Null Hypothesis |     | $I_1 = 1, I_2 = 0$ |       | $I_1 = 1, I_2 = 1$ |     |
|-----------------|-----|--------------------|-------|--------------------|-----|
|                 |     | F                  | LR    | F                  | LR  |
| Sample          | 20  | 19.2               | 28.8  | 1.3                | 2.5 |
|                 | 30  | 83.6               | 88.8  | 1.1                | 3.4 |
|                 | 40  | 97.0               | 98.4  | 1.1                | 2.0 |
| Size            | 60  | 99.9               | 99.9  | 1.1                | 1.1 |
|                 | 100 | 100.0              | 100.0 | 1.0                | 1.0 |



Table 4b

Size/Power of Test at 5% Significance Level Nominal

True Model:  $I_1=1, I_2=1$

| Null Hypothesis |     | $I_1 = 1, I_2 = 0$ |       | $I_1 = 1, I_2 = 1$ |     |
|-----------------|-----|--------------------|-------|--------------------|-----|
| Test            |     | F                  | LR    | F                  | LR  |
| Sample          | 20  | 42.7               | 44.7  | 5.7                | 6.9 |
|                 | 30  | 95.8               | 96.3  | 5.3                | 7.7 |
|                 | 40  | 99.6               | 99.8  | 4.7                | 5.7 |
| Size            | 60  | 99.9               | 100.0 | 5.2                | 5.2 |
|                 | 100 | 100.0              | 100.0 | 5.1                | 5.0 |

Table 4c

Size/Power of Test at 10% Significance Level Nominal

True Model:  $I_1=1, I_2=1$

| Null Hypothesis |     | $I_1 = 1, I_2 = 0$ |       | $I_1 = 1, I_2 = 1$ |      |
|-----------------|-----|--------------------|-------|--------------------|------|
| Test            |     | F                  | LR    | F                  | LR   |
| Sample          | 20  | 56.1               | 57.7  | 10.9               | 10.5 |
|                 | 30  | 98.2               | 98.2  | 10.2               | 10.9 |
|                 | 40  | 99.9               | 99.9  | 10.9               | 9.3  |
| Size            | 60  | 100.0              | 100.0 | 11.1               | 9.9  |
|                 | 100 | 100.0              | 100.0 | 10.1               | 9.4  |

## 6. Conclusion

We have derived an exact test for the parameter vector of transformation in linear models. By utilizing Taylor series approximations this reduces to a choice between two regression equations. The foregoing analysis need not specifically assume the transformation parameters to be 0 or 1. Our proposed test is thus applicable to any other transformation parameter values though interpretation is straightforward when they are equal to 1 or 0. The test statistic which has an exact  $F$ -distribution can be easily calculated from these two regression equations by least squares algorithm. Monte Carlo results have demonstrated that our proposed procedure is generally more capable than the likelihood approach in stating the correct size of the test, yet it is equally powerful to the latter in rejecting false null hypotheses. It is therefore a simple and ready alternative to the likelihood ratio test for assessing the suitable choice of the combination of first differences and percentage changes in linear models, thereby allowing more flexible and appropriate economic relations be formulated and their validity be tested.

## Acknowledgment

This research was supported by a grant from The Hong Kong Polytechnic University Research Committee.

## References

- Andersen, L.C. and K.M. Carlson (1970): "A monetarist model for economic stabilization," *Review of the St. Louis Federal Reserve Bank*, 52, 7-25.
- Andrews, D.F. (1971): "A note on the selection of data transformations," *Biometrika*, 58, 249-254.
- Colclough, W.G. and Lange, M.D. (1982): "Empirical evidence of causality from consumer to wholesale prices," *Journal of Econometrics*, 19, 379-384.
- Coulson, N.E. and R.P. Robins (1985): "A comment on the testing of functional form in first difference models," *Review of Economics and Statistics*, 67, 710-712.
- Box, G.E.P. (1980): "Sampling and Bayes' inference in scientific modelling and robustness (with Discussion)," *Journal of the Royal Statistical Society A*, 143, 383-430.

Dhrymes, P.J., E.P. Howrey, S.H. Hymans, J. Kmenta, E.E. Leamer, R.E. Quandt, J.B. Ramsey, H.T. Shapiro and V. Zaronwitz (1972): "Criteria for evaluation of econometric models," *Annals of Economic and Social Measurements*, 1, 291-324.

Godfrey, L.G. and M.R. Wickens (1981): "Testing linear and log-linear regressions for functional form," *Review of Economic Studies*, 48, 497-496.

Hafer, R.W. and S.E. Hein (1980): "The dynamics and estimation of short-run money demand," *Review of the St. Louis Federal Reserve Bank*, 62, 26-35.

Layson, S.K. and T.G. Seaks (1984): "Estimation and testing for functional form in first difference models," *The Review of Economics and Statistics*, 66, 338-343.

Milliken, G.A. and F.A. Graybill (1970): "Extensions of the general linear hypothesis model," *Journal of the American Statistical Association*, 65, 797-807.

Park, T. (1991): "Double length regressions for testing the Box-Cox difference transformation," *The Review of Economics and Statistics*, 73, 181-185.

Seaks, T.G. and D.P. Vines (1990): "A Monte Carlo evaluation of the Box-Cox difference transformation," *The Review of Economics of Economics and Statistics*, 72, 506-510.

Thornton, J.R., R.J. Agnello and C.R. Link (1978): "Poverty and economic growth: Trickle down peters out," *Economic Inquiry*, 16, 385-394.

Ward, G.C. and I.D. Dick (1952): "Non-additivity in randomized block designs and balanced block designs," *New Zealand Journal of Science and Technology*, 33, 430-436.